

MAGNETOTAIL SIGNATURES OF SUBSTORMS ASSOCIATED WITH SHEATH AND CIR REGIONS IN THE SOLAR WIND

Irina Despirak¹, Andris Liubchich¹, Rositza Koleva²

¹Polar Geophysical Institute – Russian Academy of Sciences, Apatity, Murmansk region, Russia

²Space Research and Technology Institute – Bulgarian Academy of Sciences
e-mail: despirak@gmail.com

Keywords: solar wind, magnetosphere, substorms

Abstract: It is well known that the various structures in the solar wind flow are either related to the solar corona structure or are created along solar wind propagation. Structures created while the solar wind propagates through the Solar system are the regions of compressed plasma at the fronts of recurrent solar wind streams (CIR) and magnetic clouds (Sheath). Auroral disturbances observed by Polar spacecraft during Sheath and CIR events had an unusual form. On one side these disturbances exhibited typical substorm signatures, but on the other side they had very large longitudinal dimensions and covered a very large area. To examine whether these disturbances represent substorms we considered data from the Geotail spacecraft in the magnetotail. Auroral bulge parameters were obtained by the UVI onboard Polar; solar wind parameters were taken from the OMNI data base. All auroral disturbances observed by Polar during Sheaths and CIRs for the periods 1997-1998; 2000; October 2001 and December 1996 were studied. We show that some signatures of a typical substorm were observed in the magnetotail, namely: 1) fast plasma flows associated with a reconnection process (tailward/earthward flows) 2) a sharp decrease of the total pressure following the interval of a pressure increase. This enables us to consider the auroral disturbances during the fronts of high-speed streams as substorm.

Introduction

Solar wind is not a uniform flow, various large-scale structures and streams exist within it [1]. The streams and structures in the solar wind could be two types: quasi-stationary and disturbed [2]. To the disturbed type belong the interplanetary manifestations of coronal mass ejections that can include magnetic clouds (MC) and EJECTA (or interplanetary coronal mass ejection). Disturbed structures are also structures generated in the interplanetary medium along solar wind propagation - compression regions in front of incoming fast recurrent streams (CIR) and magnetic clouds (Sheath) (e.g. [3], [4]). The disturbed solar wind structures – magnetic clouds (MC) and the compression regions in front of streams (Sheath and CIR), have a noticeable in magnitude and duration south component of the magnetic field ($B_z < 0$) and could trigger geomagnetic storms [5]. The relationship between storm time substorms and isolated substorms is still an open question. Many researches find no differences between storm-time and nonstorm classical substorms, while others find substantial difference between them, e.g. during storm-time substorms they observe lack of bulge, lack of bifurcation of aurora (e.g. [6]; [7]).

Recently we considered auroral bulge developments during 4 solar wind structures: magnetic clouds (MC), recurrent streams (RS), and regions of their interaction with undisturbed solar wind (Sheath and CIR) [8]. It was demonstrated that during the impact of compression regions Sheath and CIR peculiar auroral disturbances of large area and very large longitudinal and latitudinal dimensions are observed.




Fig. 1. Examples of auroral disturbances development by Polar UVI data during Sheath, 17 September 2000 (a) and during CIR, 28 February 1997 (b). On each auroral image the blue curve delimits the bulge region.

Although the auroral disturbances during Sheaths and CIRs have signatures of an auroral substorm development - localized onset and formation of the auroral bulge, a question arises if these disturbances are substorms. To answer this question we investigate magnetotail plasma dynamics in the course of the auroral bulge formation when the magnetosphere is driven by Sheaths and CIRs.

In the course of a substorm fast plasma flows in the magnetotail are observed and satellites in the near or middle tail can register a reversal of a tailward plasma flow to an earthward plasma flow. The observation of fast flow reversal by a spacecraft in the plasma sheet is a substorm signature - it indicates passage of the reconnection site around the spacecraft in the NENL model (e.g. [9]) as shown in Fig. 2, or passage of the current disruption region in the Current Disruption model (e.g. [10]).




Fig. 2. Schematic view of the situation when a satellite in the magnetotail observes a reversal of fast flows. The region where tailward and earthward flows (shown by arrows) are generated is associated with the X-line. (Figure taken from [9])

During substorm growth phase, as a result of reconnection at the magnetopause magnetic flux is stored in the magnetotail lobes, and in the course of an auroral disturbances this flux is 'unloaded' down to the ionosphere. So the total pressure increase followed by a decrease observed during the substorm-related fast flows is one of the signatures of substorm development in the magnetotail.

Data used

The auroral disturbances are studied by Polar UVI data in the LBHL band (1600-1800Å); Geotail plasma measurements by LEP instrument, and magnetic field measurements by MGF instrument; the solar wind and interplanetary magnetic field parameters measured by Wind spacecraft were taken from OMNI database.

The events were selected using the following criteria:

- 1) The auroral disturbances should be observed by the UVI onboard Polar;
- 2) The auroral disturbances should be observed during Sheath, CIR, MC or RS solar wind structures;
- 3) The meridian of the Geotail footprint should cross the auroral bulge;
- 4) Geotail should be in the night plasma sheet. The criterion $\beta > 0.1$ (β is the ratio of kinetic plasma pressure to magnetic pressure) and eye inspection of ion and electron spectra are applied for the plasma sheet identification

All auroral disturbances observed by Polar during Sheaths and CIRs for the periods 1997-1998; 2000; October 2001 and December 1996 were studied, 8 events when Geotail was in the plasma sheet during the auroral bulge formation connected to Sheath and CIR impact were selected. We present two of them.

Results

1) Auroral disturbances during CIR - 29 March 1997

A recurrent stream reached the Earth at about 16 UT on 29 March and passed away at about 04 UT on 31 March 1997 (as deduced from Wind data). The CIR was registered from about 06 UT to 16 UT. Substorm was observed at 14:37 UT.

The blue line on (a) delimits the intervals of CIR and RS, the interval of the recurrent stream is marked by blue crosshatched regions. The recurrent stream reached the Earth at about 16 UT on 29 March and passed away at about 04 UT on 31 March 1997. In front of the recurrent stream there is region of interaction with slower streams (CIR). These are regions with magnetic field and plasma compression. The CIR was registered about from 06 UT to 16 UT on 29 March. Vertical blue lines and red crosshatched regions show the times of CIR.

Substorm is observed during the CIR at 14:37 UT, the black solid lines show the onset time of the substorms registered by Polar satellite. The blue curve delimits the bulge region on each auroral image in (b). The onset latitude L_0 of the bulge was 63.3° CGLAT; the maximum latitude $L_m - 79^\circ$ CGLAT; the latitudinal size $L_{lat} \sim 16.3^\circ$ CGLAT, the longitudinal size $L_{long} \sim 98^\circ$ CGLAT, the ratio between the longitudinal and the latitudinal sizes L_{lat}/L_{long} was equal to ~ 6.01 . Keograms on bottom panel of (b) demonstrate the clear poleward expansion of the bulge.

Geotail data for the period 13 – 16 UT on 29 March 1997 are presented in (c). On 29 March 1997 at $\sim 05:00$ UT Geotail crossed the evening flank magnetopause and proceeded through the plasma sheet (PS) towards north. In the depicted in Fig. 3c Geotail was in the southern neutral sheet at GSM [-20.9, 14.6, 2.1] and at 13:30 moved to the northern PS, heading to the northern lobe. At 14:31 the s/c suddenly jumped to the southern PS, total pressure increased, which possibly is connected with a thinning of the PS. The satellite started registering fast tailward plasma flows about 14:40 UT, a flow reversal took place at $\sim 14:41$ UT and maximum of earthward flow is observed at 14:49 UT. The flow reversal is associated with a decrease of the total (magnetic plus plasma) pressure.




Fig. 3. Auroral disturbances during CIR - 29 March 1997

2) Auroral disturbances during SHEATH event - 10 October 1997

A magnetic cloud arrived at $\sim 23:8$ UT on 10 October and passed away at ~ 01 UT on 12 October 1997 (as deduced from Wind data). The Sheath was registered about from 16 UT to 23:8 UT. Substorm was observed at 21:39 UT. The format of Figure 4 is the same as that of Figure 3, the red crosshatched region in (4a) shows the time of the Sheath. On the top panel (4b), the auroral bulge development from 21:37 to 22:36 UT according to Polar UVI data is shown. The onset latitude L_0 of the bulge was 55.7° CGLAT; the maximum latitude $L_m - 73.7^\circ$ CGLAT; the latitudinal size $L_{lat} \sim 16^\circ$ CGLAT, the longitudinal size $L_{long} \sim 110^\circ$ CGLAT, the ratio between the longitudinal and the latitudinal sizes L_{lat}/L_{long} was equal to ~ 6.9 .




Fig. 4. Auroral disturbances during SHEATH event - 10 October 1997

Geotail data for the period 21-24 UT are shown on panel 4c. On 10 October 1997 Geotail encountered the night southern plasma sheet at ~11:30 UT near the Earth, and proceeded tailward and northwards. The coming of the Sheath produced at 16:12 a strong increase of the magnetic field and density in the plasma sheet. During the interval depicted in Fig. 4c Geotail was at GSM $\sim [-13 - 14, -1 - 5, -0.4 - 0]$. A fast tailward flow is registered with a maximum at 21:52 UT, a flow reversal – at 21:59 and earthward flows with a maximum at 22:03; a decrease in the total (magnetic plus plasma) pressure near the reversal is observed.

Discussion

In both cases presented here, Geotail in the magnetotail registered a reversal of a tailward plasma flow to an earthward plasma flow. The observation of oppositely directed flows is interpreted as tailward movement of the reconnection site ([11], [12]), the tailward flow being inside a plasmoid. In the near-Earth initiation scenario oppositely directed flows are also observed, but not organized by the magnetic field; their generation evoked by current disruption (CD), the region of current disruption and dipolarization proceeds progressively down the magnetotail (e.g. [10]). The earthward flow is observed in the depolarized region, the tailward flow – tailward of the CD region. So the observation of fast flow reversal by a spacecraft in the plasma sheet is a substorm signature - it indicates passage of the reconnection site around the spacecraft in the NENL model (e.g. [9]) or passage of the current disruption region in the CD model.

In the magnetotail at Geotail location a sharp decrease of the total pressure following the interval of pressure increase was observed. The total pressure decrease is associated with the time of plasma flow direction change. From the pressure balance condition, which is suggested to held across the plasma sheet and tail lobe boundary, the total pressure in the plasma sheet must be equal to the lobe magnetic pressure. The decrease below the lobe magnetic pressure characterizes that part of the magnetic energy stored in the tail, which dissipates during the substorm. As is well known during substorm growth phase, as a result of reconnection at the magnetopause magnetic flux is stored in the magnetotail lobes, the volume of the flux stored depending on solar wind parameters and interplanetary magnetic field (e.g. [13]). During auroral disturbances this flux is 'unloaded' down to the

ionosphere. However it is possible that during a substorm not all the stored flux but only a part of it is dissipated in the ionosphere. The area of the auroral bulge could be a measure of the dissipated magnetic flux [14], [15]. So the total pressure increase followed by a decrease observed during the substorm-related fast flows is one of the signatures of substorm development in the magnetotail.

Conclusions

In all eight cases we have analyzed, in the course of auroral disturbances development in the ionosphere during Sheath and CIR, in the magnetotail typical substorm signatures are observed:

- fast plasma flows associated with the reconnection process (tailward/earthward flows)
- a sharp decrease of the total pressure following the interval of pressure increase.

This enables us to consider the auroral disturbances during Sheath and CIR as substorms.

Acknowledgements. The paper was supported by the RFBR Grants 12-05-01030 and Program No 22 of the Presidium of the Russian Academy of Sciences (RAS) "Fundamental problems of the Solar system exploration". The work was also partially supported by grant DID 02/8 from the Bulgarian National Science Fund. The study is part of a joint Russian - Bulgarian Project "The influence of solar activity and solar wind streams on the magnetospheric disturbances, particle precipitations and auroral emissions" of PGI RAS and IKIT-BAS under the Fundamental Space Research Program between RAS and BAS.

References:

1. Pudovkin, M.I., Solar wind, Soros Educational Journal, 12, pp. 87-94, 1996.
2. Yermolaev, Yu.I., N.S. Nikolaeva, I.G. Lodkina, M.Yu. Yermolaev, Catalog of Large-Scale Solar Wind Phenomena during 1976– 2000, Kosmicheskie Issledovaniya, 47, pp. 99–113, 2009.
3. Klein, L.W., L.F. Burlaga, Interplanetary magnetic clouds at 1 AU, Journal of Geophysical Research, 87, pp. 613-624, 1982.
4. Balogh, A., J.T. Gosling, J.R. Jokipii, R. Kallenbach, H. Kunow, Corotating interaction region. Space Science Reviews, 89, pp. 141-411, 1999.
5. Yermolaev, Yu.I., N.S. Nikolaeva, I.G. Lodkina, M.Yu. Yermolaev, Relative occurrence rate and geoeffectiveness of large-scale types of the solar wind, Kosmicheskie Issledovaniya, 48, pp. 3–32, 2010.
6. McPherron, R.L., Hsu, T.-S., 2002. A comparison of substorms occurring during magnetic storms with those occurring during quiet times. Journal of Geophysical Research 107, 1259, doi:10.1029/2001JA002008.
7. Hoffman, R.A., J.W. Gjerloev, L.A. Frank, J.W. Sigwarth, Are there optical differences between storm-time substorms and isolated substorms? Annales Geophysicae, 28, pp. 1183–1198, 2010.
8. Despirak, I.V., A.A. Lubchich, A.G. Yahnin, B.V. Kozelov, H.K. Biernat, Development of substorm bulges during different solar wind structures, Annales Geophysicae, 27, pp. 1951-1960, 2009.
9. Yahnin, A.G., I.V. Despirak, A.A. Lubchich, B.V. Kozelov, N.P. Dmitrieva, M.A. Shukhtina, H.K. Biernat, Indirect mapping of the source of the oppositely directed fast plasma flows in the plasma sheet onto the auroral display. Annales Geophysicae 24, pp. 679–687, 2006a.
10. Lui, A. T. Y., A. Zheng, H. Reme, M.V. Dunlop, G. Gustafsson, Evaluation of substorm models with Cluster observations of plasma flow reversal in the magnetotail, Advances in Space Research, 41, pp. 1611–1618, 2008.
11. Hones, Jr., E. W., Transient phenomena in the magnetotail and their relation to substorms, Space Science Reviews, 16, pp. 617–410, 1979.
12. Runov, A., R. Nakamura, W. Baumjohann, R.A. Treumann, T.L. Zhang, M. Volwerk, Z. Vörös, A. Balogh, K.-H. Glaßmeier, B. Klecker, H. Rème, L. Kistler, Current sheet structure near magnetic X-line observed by Cluster, Geophysical Research Letters, 30, 1579, doi: 10.1029/2002GL016730, 2003.
13. Shukhtina, M.A., N.P. Dmitrieva, N.G. Popova, V.A. Sergeev, A.G. Yahnin, A.G., I.V. Despirak, Observational evidence of the loading-unloading substorm scheme, Geophysical Research Letters, 32, L17107, doi:10.1029/2005GL023779, 2005.
14. Miyashita, Y., S. Machida, K. Liou, T. Mukai, Y. Saito, C.-I. Meng, G.K. Parks, Relationship between magnetotail variations and auroral activities during substorms, Journal of Geophysical Research, 108, 1022, doi:10.1029/2001JA009175, 2003.
15. Yahnin, A.G., I.V. Despirak, A.A. Lubchich, B.V. Kozelov, N.P. Dmitrieva, M.A. Shukhtina, H.K. Biernat, Relationship between substorm auroras and processes in the near-earth magnetotail, Space Science Reviews, 122, pp. 97-106, 2006b.